

German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.



Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.

Provided by

wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.



Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.

Provided by

wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.



Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.

Provided by

wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.



Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.

Provided by

wimastergardener.org



Additional Resources

German Yellowjacket

- *Social Wasps and Bees in the Upper Midwest*, at <http://www.extension.umn.edu/garden/insects/find/wasp-and-bee-control/>

Additional Resources

German Yellowjacket

- *Social Wasps and Bees in the Upper Midwest*, at <http://www.extension.umn.edu/garden/insects/find/wasp-and-bee-control/>

Additional Resources

German Yellowjacket

- *Social Wasps and Bees in the Upper Midwest*, at <http://www.extension.umn.edu/garden/insects/find/wasp-and-bee-control/>

Additional Resources

German Yellowjacket

- *Social Wasps and Bees in the Upper Midwest*, at <http://www.extension.umn.edu/garden/insects/find/wasp-and-bee-control/>

German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.

Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.



Provided by
wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.

Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.



Provided by
wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.

Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.



Provided by
wimastergardener.org



German Yellowjacket

During late summer and fall, yellowjackets become aggressive scavengers, feeding on sugars and proteins. They frequently disrupt outside activities where food or drink is served, sometimes becoming a nuisance as early as the beginning of August. German yellowjackets nest in the ground or in wall voids and are most aggressive when the nest is disturbed.

Control

- Destroy nests early in the season (pour a soap and water solution into the entrance of ground nests or apply an insecticide in the opening).
- Decrease the chance of attracting wasps by waiting to serve food and drink until people are ready to eat and promptly putting away food when done. Keep garbage cans tightly sealed.
- Be very watchful when eating or drinking outdoors during the day. If a wasp flies to your food, wait for it to fly away.



Provided by
wimastergardener.org

